

- Statistisches Bundesamt (Hrsg.): Statistisches Jahrbuch 2005. S. 238
- VERA-Schriftenreihe: „Lebensmittel- und Nährstoffaufnahme in der BRD (1985–1989)“, Band XII, Wissenschaftlicher Fachverlag, Niederkleen, 1994
- Stunkard AJ, Harris JR, Pedersen NL, McClearn GE. The body-mass index of twins who have been reared apart. *N Engl J Med* 1990;322:1483–1487
- Rankinen T, Zuberi A, Chagnon YC, Weisnagel SJ, Argyropoulos G, Walts B, Pérusse L, Bouchard C. The human obesity gene map: the 2005 update. *Obesity (Silver Spring)*. 2006 Apr;14(4):529-644.
- Skender ML, Goodrick GK, Del Junco DJ, Reeves RS, Darnell L, Gotto A, et al. Comparison of 2-year weight loss trends in behavioral treatments of obesity: Diet, exercise, and combination interventions. *J Am Diet Assoc* 1996;96:342-6.
- Wing RR. Behavioural treatment of severe obesity. *Am J Clin Nutr* 1992;55(2 Suppl):545-551
- Garrow JS. Exercise in the treatment of obesity: a marginal contribution. *Int J Obes Relat Metab Disord* 1995a;19(Suppl 4):126-9, kein Abstract, Evidenzklasse: IV.
- Epstein LH, Coleman KJ, Myers MD. Exercise in treating obesity in children and adolescents. *Med Sci Sports Exerc* 1996a;28(4):428-35.
- Harrell JS, Johnston LF, Griggs TR, Schaefer P, Carr EG, Jr., McMurray RG, et al. An occupation based physical activity intervention program: improving fitness and decreasing obesity. *Aaohn J* 1996;44(8):377-84.
- Buemann B, Tremblay A. Effects of exercise training on abdominal obesity and related metabolic complications. *Sports Med* 1996;21(3):191-212. References Page 34 of 37
- Hauner H. Strategie der Adipositastherapie. *Der Internist* 1997;3:244-250.
- De Luis DA et al., *Ann Nutr Metab* 50: 354-360, 2006
- Lindi et al., *Diabetes* 51: 2581-2586, 2002
- Martinez et al., *J Nutr* 133: 2549-2554, 2003
- Masuo et al., *Circulation* 111: 3429-3434, 2005
- Marti et al., *Diabetes Obes Metab* 4: 428-430, 2002
- Shiwaku et al., *Int J Obes Relat Metab Disord* 27: 1028-1036, 2003
- Schon leichtes Übergewicht beeinträchtigt die Lebenserwartung, *MMW-Fortschr. Med. Nr. 51-52 / 2006* (148. Jg.), S. 28
- Elias, M. F. et al.: Obesity, diabetes and cognitive deficit: The Framingham Heart Study. In: *Neurobiol Aging.* 26, Nr. 1, 2005, S. 11-16.
- Wolf P. A. et al.: Relation of obesity to cognitive function: importance of central obesity and synergistic influence of concomitant hypertension. The Framingham Heart Study. In: *Curr Alzheimer Res.* 4, Nr. 2, 2007, S. 111-116.
- Irie F. et al.: Enhanced risk for Alzheimer disease in persons with type 2 diabetes and APOE epsilon4: the Cardiovascular Health Study Cognition Study. In: *Arch Neurol.* 65, Nr. 1, 2008, S. 83-89.
- Xu W. L. et al.: Uncontrolled diabetes increases the risk of Alzheimer's disease: a population-based cohort study. In: *Diabetologia.* 52, Nr. 6, 2009, S. 1031-1039.
- Naderali, E. K. et al.: Obesity and Alzheimer's Disease: A Link Between Body Weight and Cognitive Function in Old Age. In: *Am J Alzheimers Dis Other Demen..*
- Cyrus A. R. et al.: Brain Structure and Obesity. In: *Human Brain Mapping.*
- Am J Clin Nutr. 2009 Dec;90(6):1483-8. Epub 2009 Oct 14. The FTO gene rs9939609 obesity-risk allele and loss of control over eating. Tanofsky-Kraff M, Han JC, Anandalingam K, Shomaker LB, Columbo KM, Wolkoff LE, Kozlosky M, Elliott C, Ranzenhofer LM, Roza CA, Yanovski SZ, Yanovski JA.
- Am J Clin Nutr. 2009 Nov;90(5):1418-25. Epub 2009 Sep 2. Fat and carbohydrate intake modify the association between genetic variation in the FTO genotype and obesity. Sonestedt E, Roos C, Gullberg B, Ericson U, Würfält E, Orho-Melander M.
- Arch Intern Med. 2009 Nov 9;169(20):1897-906. APOA2, dietary fat, and body mass index: replication of a gene-diet interaction in 3 independent populations. Corella D, Peloso G, Arnett DK, Demissie S, Cupples LA, Tucker K, Lai CQ, Parnell LD, Coltell O, Lee YC, Ordovas JM.
- Arterioscler Thromb Vasc Biol. 1998 Oct;18(10):1606-10. Postprandial lipemic response is modified by the polymorphism at codon 54 of the fatty acid-binding protein 2 gene. Agren JJ, Valve R, Vidgren H, Laakso M, Uusitupa M.
- Clin Chem Lab Med. 2007;45(3):316-20. Polymorphisms in the APOA1/C3/A4/A5 gene cluster and cholesterol responsiveness to dietary change. Hubacek JA, Bohuslavova R, Skodova Z, Pitha J, Bobkova D, Poledne R.
- Clin Chim Acta. 2010 Nov 11;411(21-22):1716-22. Epub 2010 Jul 25. Effects of common FTO gene variants associated with BMI on dietary intake and physical activity in Koreans. Lee HJ, Kim IK, Kang JH, Ahn Y, Han BG, Lee JY, Song J.
- Clin Genet. 2003 Feb;63(2):109-16. The PPAR-gamma P12A polymorphism modulates the relationship between dietary fat intake and components of the metabolic syndrome: results from the Québec Family Study. Robitaille J, Després JP, Pérusse L, Vohl MC. Page 35 of 37
- Diabetes Care. 1997 Dec;20(12):1887-90. Effects of Trp64Arg mutation in the beta 3-adrenergic receptor gene on weight loss, body fat distribution, glycemic control, and insulin resistance in obese type 2 diabetic patients. Sakane N, Yoshida T, Umekawa T, Kogure A, Takakura Y, Kondo M.
- Diabetes Obes Metab. 2002 Nov;4(6):428-30. TRP64ARG polymorphism of the beta 3-adrenergic receptor gene and obesity risk: effect modification by a sedentary lifestyle. Martí A, Corbalán MS, Martínez-González MA, Martínez JA.
- Diabetes. 2002 Aug;51(8):2581-6. Association of the Pro12Ala polymorphism in the PPAR-gamma2 gene with 3-year incidence of type 2 diabetes and body weight change in the Finnish Diabetes Prevention Study. Lindi VI, Uusitupa MI, Lindström J, Louheranta A, Eriksson JG, Valle TT, Härmäläinen H, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, Laakso M, Tuomilehto J; Finnish Diabetes Prevention Study.
- Diabetologia. 2005 Aug;48(8):1503-9. Epub 2005 Jun 29. Influence of Pro12Ala peroxisome proliferator-activated receptor gamma2 polymorphism on glucose response to exercise training in type 2 diabetes. Adamo KB, Sigal RJ, Williams K, Kenny G, Prud'homme D, Tesson F.
- Eur J Intern Med. 2007 Dec;18(8):587-92. Epub 2007 Sep 10. Influence of the Trp64Arg polymorphism in the beta 3 adrenoreceptor gene on insulin resistance, adipokine response, and weight loss secondary to lifestyle modification in obese patients. de Luis DA, Gonzalez Sagrado M, Aller R, Izaola O, Conde R.
- Horm Metab Res. 2005 Feb;37(2):99-105. Influence of the PPAR-gamma2 Pro12Ala and ACE I/D polymorphisms on insulin sensitivity and training effects in healthy offspring of type 2 diabetic subjects. Østergård T, Ek J, Hamid Y, Saltin B, Pedersen OB, Hansen T, Schmitz O.
- Int J Obes (Lond). 2011 Aug;35(8):1041-9. doi: 10.1038/ijo.2010.263. Epub 2010 Dec 21. Association between fat intake, physical activity and mortality on genetic variation in FTO. Sonestedt E, Gullberg B, Ericson U, Würfält E, Hedblad B, Orho-Melander M.
- Int J Obes (Lond). 2011 May;35(5):666-75. Epub 2010 Oct 26. Association between the APOA2 promoter polymorphism and body weight in Mediterranean and Asian populations: replication of a gene-saturated fat interaction. Corella D, Tai ES, Sorlí JV, Chew SK, Coltell O, Sotos-Prieto M, García-Ríos A, Estruch R, Ordovas JM.
- Int J Obes Relat Metab Disord. 2003 Sep;27(9):1028-36. Difficulty in losing weight by behavioral intervention for women with Trp64Arg polymorphism of the beta3-adrenergic receptor gene.

Shiwaku K, Nogi A, Anuurad E, Kitajima K, Enkhmaa B, Shimono K, Yamane Y.

- *J Biol Chem.* 2001 Oct 26;276(43):39679-84. Epub 2001 Aug 3. The polymorphism at codon 54 of the FABP2 gene increases fat absorption in human intestinal explants. Levy E, Ménard D, Delvin E, Stan S, Mitchell G, Lambert M, Ziv E, Feoli-Fonseca JC, Seidman E.
- *J Clin Endocrinol Metab.* 1998 Jul;83(7):2441-4. Meta-analysis of the association of Trp64Arg polymorphism of beta 3-adrenergic receptor gene with body mass index. Fujisawa T, Ikegami H, Kawaguchi Y, Ogihara T.
- *J Lipid Res.* 2000 Dec;41(12):2002-8. Effects of an Ala54Thr polymorphism in the intestinal fatty acidbinding protein on responses to dietary fat in humans. Pratley RE, Baier L, Pan DA, Salbe AD, Storlien L, Ravussin E, Bogardus C.
- *J Mol Med (Berl).* 2007 Feb;85(2):119-28. Epub 2007 Jan 9. APOA5 gene variation modulates the effects of dietary fat intake on body mass index and obesity risk in the Framingham Heart Study. Corella D, Lai CQ, Demissie S, Cupples LA, Manning AK, Tucker KL, Ordovas JM. Page 36 of 37
- *J Nutr.* 2003 Aug;133(8):2549-54. Obesity risk is associated with carbohydrate intake in women carrying the Gln27Glu beta2-adrenoceptor polymorphism. Martínez JA, Corbalán MS, Sánchez-Villegas A, Forga L, Martí A, Martínez-González MA.
- *J Nutr.* 2009 Dec;139(12):2301-8. Epub 2009 Oct 14. Apolipoprotein A5 polymorphisms interact with total dietary fat intake in association with markers of metabolic syndrome in Puerto Rican older adults. Mattei J, Demissie S, Tucker KL, Ordovas JM.
- *J Nutr.* 2011 Dec;141(12):2219-25. Epub 2011 Nov 2. A High Intake of Saturated Fatty Acids Strengthens the Association between the Fat Mass and Obesity-Associated Gene and BMI. Corella D, Arnett DK, Tucker KL, Kabagambe EK, Tsai M, Parnell LD, Lai CQ, Lee YC, Warodomwichit D, Hopkins PN, Ordovas JM.
- *J Nutr.* 2011 Mar;141(3):380-5. Epub 2011 Jan 5. APOA5 gene variation interacts with dietary fat intake to modulate obesity and circulating triglycerides in a Mediterranean population. Sánchez-Moreno C, Ordovás JM, Smith CE, Baraza JC, Lee YC, Garaulet M.
- *Metabolism.* 2005 Dec;54(12):1652-8. Comparison of the acute response to meals enriched with cis or trans-fatty acids on glucose and lipids in overweight individuals with differing FABP2 genotypes. Lefèvre M, Lovejoy JC, Smith SR, Delany JP, Champagne C, Most MM, Denkins Y, de Jonge L, Rood J, Bray GA.
- *Obes Res.* 2001 Dec;9(12):741-5. Association of BMI with the beta3-adrenergic receptor gene polymorphism in Japanese: meta-analysis. Kurokawa N, Nakai K, Kameo S, Liu ZM, Satoh H.
- *Obes Res.* 2004 Feb;12(2):340-5. Intestinal FABP2 A54T polymorphism: association with insulin resistance and obesity in women. Albala C, Santos JL, Cifuentes M, Villarroel AC, Lera L, Liberman C, Angel B, Pérez-Bravo F.
- *Obes Res.* 2004 May;12(5):807-15. Association between body fat response to exercise training and multilocus ADR genotypes. Phares DA, Halverstadt AA, Shuldiner AR, Ferrell RE, Douglass LW, Ryan AS, Goldberg AP, Hagberg JM.
- *Obesity (Silver Spring).* 2006 Apr;14(4):529-64. The human obesity gene map: the 2005 update. Rankinen T, Zuberi A, Chagnon YC, Weisnagel SJ, Argyropoulos G, Walts B, Pérusse L, Bouchard C.